

## Bibliografia

### Biostimolanti

**Atiyeh R.M.**, Lee S., Edwards C.A., Arancon N.Q., Metzger J.D. 2002. The influence of humic acids derived from earthworm-processed organic wastes on plant growth. *Biores. Technol.* 84: 7-14.

**Cavani L.**, Ciavatta C., Gessa C. 2003. Determination of free L- and D-alanine in hydrolysed protein fertilisers by capillary electrophoresis. *Journal of Chromatography A*, 985: 463-469.

**Gherardi A.**, Brunelli R., Bugiani., 2000. Criteri alternativi alla “regola dei tre dieci” per la previsione delle infezioni primarie di *Plasmopara viticola*. *Atti Giornate Fitopatologiche*, vol 2, 119 - 120.

**Lachhab N.**, Sanzani S. M., Adrian M., Chiltz A., Balacey S., Boselli M., Ippolito A., P. Benoit 2014. Soybean and casein hydrolysates induce grapevine immune responses and resistance against *Plasmopara viticola*. *Front. Plant Sci.*, 5, 716.

**Martin J.**, 2012. *Impact of marine extratcs applications on cv Syrah grape (Vitis vinifera L.) yield components, harvest juice quality parameters, and nutrient uptake*. Tesi di master.

**Mugnai S.**, Azzarello E., Pandolfi C., Salamagne S., Briand X., Mancuso S., 2008. *Enhancement of ammonium and potassium root influxes by the application of marine bioactive substances positively affects Vitis vinifera plant growth*. *J. Appl. Phycol.* 20:177-182.

**Nardi, S.**, Concheri, G., Pizzeghello, D., Sturaro, A., Rella, R., Parvoli, G. 2000. Soil organic matter mobilization by root exudates. *Chemosphere* 41: 653-658.

**Paradikovic N.**, Vinkovic T., Vrcek I., Zuntar I., Bojic M., Medic-Saric M., 2011. *Effect of natural biotimulants on yield and nutritional quality: an example of sweet yellow pepper (Capsicum annuum L.) plants*. *J.Sci. Food. Agric.* 91: 2146-52.

**M. Scannavini, M.**; Spada, G.; Mazzini, E.; Cavallini, G.; Barani, A.; Ponti, I. 2000. Studio pluriennale condotto in Emilia-Romagna per la valutazione dell'efficacia di nuovi principi attivi nei confronti di *Plasmopara viticola*. *Atti Giornate Fitopatologiche*, vol 2, 169 - 176.

[www.bio\\_stimulants.eu](http://www.bio_stimulants.eu)

### Agenti di Biocontrollo

**Chet I**, Inbar J (1994) Biological control of fungal pathogens. *Appl Biochem Biotechnol* 48:37-43

**Druzhinina, I.S et al.** 2011 *Trichoderma*: the genomics of opportunistic success. *Nature Reviews*, Vol.9:749-759.

**Gullino M** (1992) Chemical control of *Botrytis* spp. In: Verhoeff K, Malathrakis NE and Williamson B (eds) *Recent Advances in Botrytis Research* (pp 217–222) Pudoc Scientific Publishers, Wageningen

**Harman G.E.**, Petzoldt R., Comis A., Chen J. (2004). Interaction between *Trichoderma harzianum* strain T22 and maize inbred line Mo 17 and effect of this interaction on disease caused by *Phytium ultimum* and *Colletotrichum graminicola*. *Phytopathology*. 94:147-153.

Hermosa, R et al. 2012 Plant-beneficial effects of *Trichoderma* and of its genes. *Microbiology*, 158, 17-25

**Howell C.R.** (2003) Mechanisms employed by *Trichoderma* species in the biological control of plant disease: the history and evolution of current concepts. *Plant Disease* 87: 4- 10.

**Klein, D.**, and Eveleigh, D. E. 1998. Ecology of *Trichoderma*. Pages 5774 in: *Trichoderma and Gliocladium*. Vol. 1. C. P. Kubicek and G. E. Harman, eds. Taylor & Francis, London. 57.

**Kubicek, C. P.**, and Harman, G. E. 1998. *Trichoderma and Gliocladium*. Vol. 1. Basic Biology, Taxonomy and Genetics. Taylor & Francis, London.

- Kubicek C.P.**, Herrera-Estrella A, Seidl-Seiboth V, *et al.* 2001 Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of *Trichoderma*.
- Shoresh M.**, Harman G.E., Mastouri F. (2010) Induced systemic resistance and plant responses to fungal biocontrol agents. *Annual Review of Phytopathology* 48:21-43
- Stanke, M *et al.* 2004 AUGUSTUS: a web server for gene finding in eukaryotes. *Nucleic Acids Research*, Vol. 32, Web Server issue W309–W312
- Sivasithamparam, K.**, and Ghisalberti, E. L. 1998. Secondary metabolism in *Trichoderma* and *Gliocladium*. Pages 139-191 in: *Trichoderma and Gliocladium*, Vol. 1. C. P. Kubicek and G. E. Harman, eds. Taylor and Francis, London.
- Woo, S.L.**, Scala, F., Ruocco, M. & Lorito, M. 2006: The molecular biology in the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology* 96, 181-185.
- Verma M.**, Brar S. K., Tyagi R.D., Surampalli R. Y., Valéro J.R. (2007) Antagonistic fungi, *Trichoderma* spp.: Panoply of biological control. *Review of Biochemical Engineering Journal* 37 (2007) 1–20
- Vinale F.**, Sivasithamparam K., Ghisalberti E.L., Marra R., Woo S.L., Lorito M. (2008) *Trichoderma* plant pathogen interactions. *Soil Biology & Biochemistry* 40: 1–10.

### Corroboranti

- Cravero S.**, Bassignana E., Crovella P., Ferrari D. (2002), Difesa della Peronospora della vite in viticoltura biologica: Verifica dell'attività fungicida di prodotti di natura diversa impiegati in vigneti piemontesi, *Atti giornate fitopatologiche 2002*, (2), pp. 289-296.
- Donnarumma L.**, La Torre A., Acciarino C. (1999), La protezione delle colture in agricoltura biologica, in *Notiziario ERSA*, (2), pp. 40-45.
- Quarles W.** (2004), Least-toxic controls of plant diseases, online, <http://www.bbg.org/gar2/topics/sustainable/handbooks/naturaldisease/leasttoxic.html>.
- Schenk P.**, Anton I., Fluri P. (2004), Effetto dell'olio di margosa sugli acari della varroa e sulle api, online, [http://www.apis.admin.ch/it/krankheiten/docs/aetherisch/niemoel\\_i.pdf](http://www.apis.admin.ch/it/krankheiten/docs/aetherisch/niemoel_i.pdf)
- Tamm L.**, Fuchs J.G., Boger N., Muhletaler L., Amsler A., levite D., Raseli A. (2004), Properties of acidified clay preparation. Eigenschaften von tonerdepreparaten: Erfahrungen aus der schweiz. Betrag am 1, Internationalen Symposium für ökologischen Weinbau. Intervist Stuttgart, 12-13 Mai 2004, pp. 27-36.

### Induttori di Resistenza

- Akiyama K.**, Kawazu K., Kobayashi A. (1995): Partially N-deacetylated chitin oligomers (pentamer to heptamer) are potential elicitors for (+)-pisatin induction in pea epicotyls, «*Zeitschrift für Naturforschung C*», 50, pp. 391-397.
- Aranega-Bou P.**, de la O Leyva M., Finiti I., García-Agustín P., González-Bosch C. (2014): Priming of plant resistance by natural compounds. Hexanoic acid as a model, «*Frontiers in Plant Science*», 5, art. 488, DOI 10.3389/fpls.2014.00488.
- Aziz A.**, Gauthier A., Bézier A., Poinssot B., Joubert J.M., Pugin A., Heyraud A., Baillieul F. (2007): Elicitor and resistance-inducing activities of beta-1,4 cellodextrins in grapevine, comparison with beta-1,3 glucans and alpha-1,4 oligogalacturonides, «*Journal of Experimental Botany*», 58, pp. 1463-1472.

- Boller T.**, Felix G. (2009): A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors, «Annual Review of Plant Biology», 60, pp. 379-406.
- Bueter C.L.**, Specht C.A., Levitz S.M. (2013): Innate sensing of chitin and chitosan, «PLoS Pathogens», 9, e1003080, DOI 10.1371/journal.ppat.1003080.
- Burketova L.**, Trda L., Ott P.G., Valentova O. (2015): Bio-based resistance inducers for sustainable plant protection against pathogens, «Biotechnology Advances», 33, pp. 994-1004.
- Chujo T.**, Takai R., Akimoto-Tomiya C., Ando S., Minami E., Nagamura Y., Kakuc H., Shibuyac N., Yasudad M., Nakashitad H., Umemurae K., Okadaa A., Okadaa K., Nojiria H., Yamanea H. (2007): Involvement of the elicitor-induced gene OsWRKY53 in the expression of defense-related genes in rice, «Biochimica et Biophysica Acta», 1769, pp. 497-505.
- Faoro F.**, Gozzo F. (2015): Is modulating virus virulence by induced systemic resistance realistic?, «Plant Science», 234, pp. 1-13.
- Gao Q.-M.**, Kachroo A., Kachroo P. (2014): Chemical inducers of systemic immunity in plants, «Journal of Experimental Botany», 65, pp. 1849-1855. Gao Q.-M., Zhu S., Kachroo P., Kachroo A. (2015): Signal regulators of systemic acquired resistance, «Frontiers in Plant Science», 6, art. 228, DOI 10.3389/fpls.2015.00228
- Gozzo F.**, Faoro F. (2013): Systemic acquired resistance (50 years after discovery): Moving from the lab to the field, «Journal of Agricultural and Food Chemistry», 61, pp. 1247312491.
- Graham M.Y.**, Weidner J., Wheeler K., Pelow M.J., Graham T.L. (2003): Induced expression of pathogenesis-related protein genes in soybean by wounding and the *Phytophthora sojae* cell wall glucan elicitor, «Physiological and Molecular Plant Pathology», 63, pp. 141-149.
- Iriti M.**, Faoro F. (2009): Chitosan as a MAMP, searching for a PRR, «Plant Signaling and Behavior», 4, pp. 66-68.
- Köhle H.**, Jeblick W., Poten F., Blaschek W., Kauss H. (1985): Chitosan-elicited callose synthesis in soybean cells as a Ca<sup>2+</sup>-dependent process, «Plant Physiology», 77, pp. 544-551.
- Landi L.**, Feliziani E., Romanazzi G. (2014): Expression of defense genes in strawberry fruits treated with different resistance inducers, «Journal of Agricultural and Food Chemistry», 62, pp. 3047-3056.
- Li Y.**, Yin H., Wang Q., Zhao X., Du Y., Li F. (2009): Oligochitosan induced *Brassica napus* L. production of NO and H<sub>2</sub>O<sub>2</sub> and their physiological function, «Carbohydrate Polymers», 75, pp. 612-617.
- Nita-Lazar M.**, Heyraud A., Gey C., Braccini I., Lienart Y. (2004): Novel oligosaccharides isolated from *Fusarium oxysporum* L. rapidly induce PAL activity in *Rubus* cells, «Acta Biochimica Polonica», 51, pp. 625-634.
- Romanazzi G.**, Murolo S., Feliziani E. (2013): A new approach to manage phytoplasma diseases: field treatments with resistance inducers to contain grapevine Bois noir, «Phytopathology», 103, pp. 785-791.
- Schreiber K.**, Desveaux D. (2008): Message in a bottle: Chemical biology of induced disease resistance in plants, «The Plant Pathology Journal», 24, pp. 245-268.
- Shah J.**, Chaturvedi R., Chowdhury Z., Venables B., Petros R.A. (2014): Signaling by small metabolites in systemic acquired resistance, «The Plant Journal», 79, pp. 645658.

**Shinya T.**, Ménard R., Kozone I., Matsuoka H., Shibuya N., Kauffmann S., Matsuoka K., Saito M. (2006): Novel  $\beta$ -1,3-,1,6-oligoglucan elicitor from *Alternaria alternata* 102 for defense responses in tobacco, «The FEBS Journal», 273, pp. 2421-2431.

**Sivakumar D.**, Bautista-Baños S. (2014): A review on the use of essential oils for postharvest decay control and maintenance of fruit quality during storage, «Crop Protection», 64, pp. 27-37.

**Stadnik M.J.**, de Freitas M.B. (2014): Algal polysaccharides as source of plant resistance inducers, «Tropical Plant Pathology», 39, pp. 111-118.

**Vera J.**, Castro J., Contreras R.A., González A., Moenne A. (2012): Oligo-carrageenans induce a long-term and broad-range protection against pathogens in tobacco plants (var. Xanthi), «Physiological and Molecular Plant Pathology», 79, pp. 31-39.

**Walters D.R.**, Ratsep J., Havis N.D. (2013): Controlling crop diseases using induced resistance: challenges for the future, «Journal of Experimental Botany», 64, pp. 12631280.

**Wolski E.A.**, Maldonado S., Daleo G.R., Andreu A.B. (2006): A novel  $\alpha$ -1, 3-glucan elicits plant defense responses in potato and induces protection against *Rhizoctonia solani* AG-3 and *Fusarium solani* f.sp. eumartii, «Physiological and Molecular Plant Pathology», 69, pp. 93-103.

**Zuppini A.**, Baldan B., Million R., Favaron F., Navazio L., Mariani P. (2004): Chitosan induces Ca<sup>2+</sup>-mediated programmed cell death in soybean cells, «New Phytologist», 161, pp. 557-568.

## **Biodiversità microbica**

Fierer, N., 2017. Embracing the unknown: disentangling the complexities of the soil microbiome. *Nature Review Microbiology* 15, 579–590. <https://doi.org/10.1038/nrmicro.2017.87> .

Veresoglou, S., Halley, J., & Rillig, M. 2015. Extinction risk of soil biota. *Nature Communications*, 6.

Mocali, S., Kuramae, E. E., Kowalchuk, G. A., Fornasier, F., Priori, S. 2020. Microbial functional diversity in vineyard soils: sulfur metabolism and links with grapevine plants and wine quality. *Frontiers in Environmental Science*, 8, 75

## **Mesofauna**

Barker K.R., Koenning S.R., 1998. Developing sustainable systems for nematode management. *Annu. Rev. Phytopathol.*, 36: 165-205.

Bongers T, 1990. The maturity index – an ecological measure of environmental disturbance based on nematode species composition. *Oecologia*, 83: 14-19.

Giller KE, Beare MH, Lavelle P., A-MN Izac, Swift MJ, 1997. Agricultural intensification, soil biodiversity and agroecosystem function. *Applied Soil Ecology*, 6(1): 3-16.

Jacomini, C., Nappi, P., Sbrilli, G., Mancini, L., 2000. Indicatori ed Indici eco tossicologici e biologici applicati al suolo: stato dell'arte. Agenzia Nazionale per la Protezione dell'Ambiente (ANPA).

Parisi V, Menta C, Gardi C, Jacomini C, Mozzanica E, 2005. Microarthropod communities as a tool to assess soil quality and biodiversity: a new approach in Italy. *Agr. Ecosys. Environ.*, 106: 323-333.

Salome C., Coll P., Lardo E., Villenave C., Blanchard E., Hinsinger P., Marsden C., Le Cadre E., 2014. Relevance of use-invariant soil properties to assess soil quality of vulnerable ecosystems: The case of Mediterranean vineyards. *Ecological Indicators*, 43: 83-93.

## **I protocolli di gestione Green Grapes del vigneto**

Arora, S., Mogha, N. K., Bhardwaj, T., Sahu, V., & Dubey, S. C. (2015). A biorational approach for management of key plant pathogens. *Indian Journal of Agricultural Sciences*, 85(6), 759-63.

Akanmu, A. O., Babalola, O. O., Venturi, V., Ayilara, M. S., Adeleke, B. S., Amoo, A. E., ... & Glick, B. R. (2021). Plant disease management: Leveraging on the plant-microbe-soil interface in the biorational use of organic amendments. *Frontiers in Plant Science*, 1590.

Horowitz, A. R., Ellsworth, P. C., & Ishaaya, I. (2009). Biorational pest control—an overview. *Biorational Control of Arthropod Pests*, 1-20.

Baysal-Gurel, F., & Bika, R. (2021). Management of Powdery Mildew on Ninebark Using Sanitizers, Biorational Products, and Fungicides. *HortScience*, 56(5), 532-537.